

On The Hermite Based-Second Kind Genocchi Polynomials

Burak Kurt and Yilmaz Simsek

Department of Mathematics, Faculty of Science University of AkdenizTR-07058 Antalya,
Turkey burakkurt@akdeniz.edu.tr and ysimsek@akdeniz.edu.tr

The 24th Congress of the Jangjeon Mathematical Society 21 July
Konya-TURKEY
Abstract

The aim of this paper is to study generating function of the Hermite-Kampé de Fériet based second kind Genocchi polynomials. We also give some identities related to these polynomials.

2010 Mathematics Subject Classification. 05A10, 11B65, 11B68.

Key Words and Phrases. Genocchi polynomials, Genocchi numbers, Second kind Bernoulli polynomials, Second kind Euler polynomials, Second kind Genocchi polynomials, Hermite polynomials, two variable Hermite polynomials, Hermite based-second kind Genocchi polynomials.

1. INTRODUCTION, DEFINITIONS AND NOTATIONS

The Bernoulli numbers the Bernoulli polynomials, the Euler numbers, the Euler polynomials and Genocchi polynomials and numbers are used many areas of sciences. These numbers and polynomials are also used in calculus of finite differences, in numerical analysis, in analytical number theory.

Recently many authors have studied on these numbers and polynomials. They also defined many different generating functions of these polynomials and numbers.

Due to Bretti and Ricci [1], the Hermite-Kampé de Fériet (or Gould-Hopper) polynomials have been used in order to construct addition formulas for different classes of generalized Gegenbauer polynomials. These polynomials (two-variable Hermite polynomials) are defined by means of the following generating functions:

$$e^{xt+yt^j} = \sum_{n=0}^{\infty} H_n^{(j)}(x, y) \frac{t^n}{n!}. \quad (1.1)$$

From (1.1), one can easily see that

$$H_n^{(j)}(x, y) = n! \sum_{r=0}^{\lfloor \frac{n}{j} \rfloor} \frac{x^{n-jr} y^r}{r! (n-jr)!},$$

where $j \geq 2$ is an integer cf. [1]. In [1], the case $j = 1$ is not considered, since the corresponding $2D$ polynomials are simply expressed by the Newton binomial formula.

The polynomials $H_n^{(j)}(x, y)$ are the solution of the generalized heat equation:

$$\begin{aligned} \frac{\partial}{\partial x} F(x, y) &= \frac{\partial^j}{\partial x^j} F(x, y), \\ F(x, 0) &= x^n. \end{aligned}$$

The classical Genocchi numbers G_n and the classical Genocchi numbers $G_n(x)$ are usually defined by means of the following generating function,

$$g(t) = \frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \quad |t| < \pi \quad (1.2)$$

and

$$g(t, x) = g(t)e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}, \quad |t| < \pi. \quad (1.3)$$

From (1.2) and (1.3), one can easily see that

$$G_n(x) = \sum_{k=0}^n \binom{n}{k} G_k x^{n-k}.$$

The second kind Genocchi polynomials of higher order are defined by means of the following generating function:

$$\mathfrak{g}(t, x) = \left(\frac{2t}{e^t + e^{-t}} \right)^a e^{tx} = \sum_{n=0}^{\infty} \mathcal{G}_n^{(a)}(x) \frac{t^n}{n!}, \quad |t| < \frac{\pi}{2}. \quad (1.4)$$

Note that $\mathcal{G}_n^{(1)}(x) = \mathcal{G}_n(x)$ denotes the second kind Genocchi polynomials cf. [15]. $\mathcal{G}_n(0) = \mathcal{G}_n$ which denote so-called the second kind Genocchi numbers cf. [15].

By using (1.4), one can easily see that

$$\mathcal{G}_n(x) = \sum_{k=0}^n \binom{n}{k} x^{n-k} \mathcal{G}_k.$$

By (1.3) and (1.4), we have

$$\begin{aligned} \mathcal{G}_n(x) &= 2^{n-1} G_n\left(\frac{x+1}{2}\right) \\ &= \sum_{k=0}^n \sum_{j=0}^{n-k} \binom{n}{k, j, n-k-j} 2^{k-1} x^{n-k-j} G_k, \end{aligned}$$

where

$$\binom{n}{a, b, c} = \frac{n!}{a!b!c!}, \quad a + b + c = n.$$

The second kind Euler polynomials of higher order are defined by means of the following generating function:

$$\left(\frac{2}{e^t + e^{-t}}\right)^a e^{xt} = \sum_{n=0}^{\infty} \mathcal{E}_n^{(a)}(x) \frac{t^n}{n!}, \quad |t| < \frac{\pi}{2} \quad (1.5)$$

Observe that $\mathcal{E}_n^{(1)}(x) = \mathcal{E}_n(x)$ denotes the second kind Euler polynomials cf. ([12], [16]).

Relation between $\mathcal{E}_n(x)$ and $\mathcal{G}_n(x)$ is given by

$$\mathcal{G}_n(x) = n\mathcal{E}_{n-1}(x).$$

Lemma 1. *The second kind Genocchi polynomials of α order is satisfied the following relations*

$$G_n^{(\alpha+\beta)}(x+x_1) = \sum_{k=0}^n \binom{n}{k} G_k^{(\alpha)}(x) G_{n-k}^{(\beta)}(x_1). \quad (1.6)$$

Also

$$\sum_{k=0}^n \binom{n}{k} E_k^{(\alpha)} G_{n-k}^{(\alpha)}(x+x_1) = \sum_{k=0}^n \binom{n}{k} E_k^{(\alpha)}(x) G_{n-k}^{(\alpha)}(x_1). \quad (1.7)$$

Proof. From (1.4),

$$\begin{aligned} \sum_{n=0}^{\infty} G_n^{(\alpha+\beta)}(x+x_1) \frac{t^n}{n!} &= \left(\sum_{n=0}^{\infty} G_n^{(\alpha)}(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} G_n^{(\beta)}(x_1) \frac{t^n}{n!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} G_k^{(\alpha)}(x) G_{n-k}^{(\beta)}(x_1) \right) \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients, we prove Lemma 1. \square

2. HERMITE-KAMPÉ DE FÉRIET BASED SECOND KIND GENOCCHI POLYNOMIALS

The aim of this section is to define the Hermite-Kampé de Fériet (or Gould-Hopper) based second kind Genocchi polynomials. Some properties of these polynomials are also given.

The Hermite-Kampé de Fériet based second kind Genocchi polynomials of higher order are defined by means of the following generating function:

$$F_{H,a}(t, x, y; j) = \left(\frac{2t}{e^t + e^{-t}}\right)^a e^{xt+yt^j} = \sum_{n=0}^{\infty} ({}_H\mathcal{G}_n^{(j,a)}(x, y)) \frac{t^n}{n!}. \quad (2.1)$$

From (2.1), one finds that

$${}_H\mathcal{E}_{n-1}^{(j,a)}(x, y) = \frac{{}_H\mathcal{G}_n^{(j,a)}(x, y)}{n},$$

where ${}_H\mathcal{E}_{n-1}^{(j,a)}(x, y)$ denotes the Hermite-Kampé de Fériet based second kind Euler polynomials of higher order.

Now we compute the derivative of ((2.1) with respect to x to derive a derivative formula for the Hermite-Kampé de Fériet based second kind Genocchi polynomials of higher order:

$$\frac{\partial}{\partial x} ({}_H\mathcal{G}_n^{(j,a)}(x, y)) = n ({}_H\mathcal{G}_{n-1}^{(j,a)}(x, y)).$$

If we substitute $j = 2$ and $a = 1$ into (2.1), and using (1.4), we have

$$\sum_{n=0}^{\infty} ({}_H\mathcal{G}_n^{(2,1)}(x, y)) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \mathcal{G}_n(x) \frac{t^n}{n!} \sum_{n=0}^{\infty} \frac{y^n t^{2n}}{n!}. \quad (2.2)$$

After some elementary calculations in the above, we obtain

$$\sum_{n=0}^{\infty} ({}_H\mathcal{G}_n^{(2)}(x, y)) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left(n! \sum_{l=0}^{\lfloor \frac{n}{2} \rfloor} \frac{y^l \mathcal{G}_{n-2l}(x)}{l!(n-2l)!} \right) \frac{t^n}{n!}.$$

By comparing coefficients of t^n in the both sides of the above, we arrive at the result of the theorem:

Theorem 1.

$${}_H\mathcal{G}_n^{(2,1)}(x, y) = n! \sum_{l=0}^{\lfloor \frac{n}{2} \rfloor} \frac{y^l \mathcal{G}_{n-2l}(x)}{l!(n-2l)!}.$$

Theorem 2.

$${}_H\mathcal{G}_n^{(j,a+b)}(x_1 + x_2, y_1 + y_2) = \sum_{k=0}^n \binom{n}{k} ({}_H\mathcal{G}_n^{(j,a)}(x_1, y_1)) ({}_H\mathcal{G}_{n-k}^{(j,b)}(x_2, y_2)). \quad (2.3)$$

Proof. By (2.1), we define the following functional equation

$$F_{H,a+b}(t, x_1 + x_2, y_1 + y_2; j) = F_{H,a}(t, x_1, y_1; j) F_{H,b}(t, x_2, y_2; j)$$

By using the above functional equation, we obtain

$$\sum_{n=0}^{\infty} ({}_H\mathcal{G}_n^{(j,a+b)}(x_1 + x_2, y_1 + y_2)) \frac{t^n}{n!} = \sum_{n=0}^{\infty} ({}_H\mathcal{G}_n^{(j,a)}(x_1, y_1)) \frac{t^n}{n!} \sum_{n=0}^{\infty} ({}_H\mathcal{G}_n^{(j,b)}(x_2, y_2)) \frac{t^n}{n!}$$

By using Cauchy product in the above equation, than comparing the coefficients of $\frac{t^n}{n!}$ on both sides, we arrive at the desired result. \square

Relation between Hermite-based second kind Genocchi polynomials of higher order, the second kind Genocchi polynomials of higher order and two-variable generalized Hermite polynomials is given by the next theorem.

Theorem 3.

$${}_H\mathcal{G}_n^{(2,a)}(x, y) = \sum_{l=0}^n \binom{n}{l} \mathcal{G}_{n-l}^{(a)} H_l^{(2)}(x, y).$$

Proof. If we substitute $j = 2$ and $a = 1$ into (2.1), we easily arrive at the desired result. \square

Acknowledgement 1. *The present investigation was supported, by the Scientific Research Project Administration of Akdeniz University.*

REFERENCES

- [1] G. Bretti and P. E. Ricci, Multidimensional extensions of the Bernoulli and Appell polynomials, Taiwanese J. Math. 8(3) (2004), 415-428.
- [2] G. Dattoli, S. Lorenzutta and C. Cesarano, Finite sums and generalization forms of Bernoulli polynomials, Rend. di Matematica, 19 (1999) 385-391.
- [3] G. Dattoli, S. Lorenzutta and C. Cesarano, Bernoulli numbers and polynomials from a more general point of view, Rend. di Matematica, 22 (2002) 193-202.
- [4] A. F. Horadam, Negative order Genocchi polynomials. Fibonacci Quart. 30 (1992), no. 1, 21-34.
- [5] S. Khan, G. Yasmin, R. Kahan and N. A. Makboul Hassan, Hermite-based Appell polynomials: properties and applications. J. Math. Anal. Appl. 351 2009(2), 756-764.
- [6] T. Kim, On the q -extension of Euler and Genocchi numbers. J. Math. Anal. Appl. 326 (2007), no. 2, 1458-1465.
- [7] T. Kim, Note on q -Genocchi numbers and polynomials. Adv. Stud. Contemp. Math. 17 (2008), no. 1, 9-15.
- [8] T. Kim, On the multiple q -Genocchi and Euler numbers. Russ. J. Math. Phys. 15 (2008), no. 4, 481-486.
- [9] T. Kim, L.-C. Jang and H. K. Pak, A note on q -Euler and Genocchi numbers. Proc. Japan Acad. Ser. A Math. Sci. 77 (2001), no. 8, 139-141.
- [10] D.-Q. Lu, Some properties of Bernoulli polynomials and their generalizations, Appl. Math. Lett. 24 (2011), no. 5, 746-751.
- [11] D.-Q. Lu, L.-C. Jang, K.-W. Hwang and Y.-Hee Kim, A note on (h, q) -Genocchi polynomials and numbers of higher order. Advnces in Dif. Equa. vol.2010, doi:10.1155/2010/309480.
- [12] L.-C. Jang and T. Kim, A new approach to q -Euler numbers and polynomials. J. Concr. Appl. Math. 6 (2008), no. 2, 159-168.
- [13] H. Ozden, Y. Simsek and H. M. Srivastava, A unified presentation of the generating functions of the generalized Bernoulli, Euler and Genocchi polynomials, Comput. Math. Appl. 60(2010), no.10 2779-2787, [DOI: 10.1016/j.camwa.2010.09.031].
- [14] S.-H. Rim, K. H. Park and E. J. Moon, On Genocchi numbers and polynomials, Abstr. Appl. Anal. 2008, Art. ID 898471, 7 pp. doi:10.155/2008/898471.
- [15] C.-S. Ryoo, A note on the second kind Genocchi polynomials, J. Comp. Anal. Appl. 13(6) (2011), 1012-1018.
- [16] C.-S. Ryoo, A note on the q -Hurwitz Euler zeta functions, J. Comp. Anal. Appl. 13(5) (2011), 986-992.
- [17] C.-S. Ryoo and T. Kim, Calculating zeros of the twisted q -Genocchi polynomials. Proc. Jangjeon Math. Soc. 12 (2009), no. 3, 343-353.
- [18] Y. Simsek, q -analogue of twisted l -series and q -twisted Euler numbers. J. of Number Theory, 110 (2005), 267-278.
- [19] Y. Simsek, I. N. Cangül, V. Kurt and D. Kim, q -Genocchi numbers and polynomials associated with q -Genocchi type l -functions. Adv. in dif. Equa. vol. 2008, doi:10.1155/2008/815750.
- [20] H. M. Srivastava, Some formulas for the Bernoulli and Euler polynomials at rational arguments. Math. Proc. Cambridge Philos. Soc. 129 (2000)(1), 77-84.
- [21] H. M. Srivastava, M. Garg and S. Choudhary, Some new families of the generalized Euler and Genocchi polynomials, Taiwanese J. Math. 15 (2011) 283-305.